

Chest Pain

Aspirin, 162-324 mg. (2-4 tabs 81mg PO) if not already administered.

Nitroglycerin (NTG), 1 tab or spray SL, If BP > 100mm/Hg systolic. May repeat every 5 min x2, if BP >100mm/Hg systolic, to a max of three doses, including any doses the patient may have self administered prior to EMS arrival.

Asthma/COPD

6mo - 2y/o: **Albuterol Sulfate** 1.25 mg
2y/o & up: **Albuterol Sulfate** 2.5 mg - 3.0 mg

Contraindications:

No DX of RAD (asthma, COPD)

History of Cardiac Disease (MI, Angina)

No current RX for fast acting beta agonists MDI/Neb

Eligible RX's: **albuterol sulfate** (Airet, Proventil, Ventolin, ProAir), **bitolterol mesylate** (Tornalate), **isoetharine** (Bronkometer, Bronkosol), **isoproterenol hydrochloride** (Isuprel), **metaproterenol sulfate** (Alupent, Metaprel), **pirbuterol acetate** (Maxair), other beta agonists

Allergic Reaction / Anaphylaxis

Severe Distress: Stridor, bronchospasm, severe abdominal pain, respiratory distress, tachycardia, shock (systolic BLOOD PRESSURE <90), observe for edema of lips, tongue or face and generalized urticaria.

PT MUST be Between 5 y/o - 65 y/o

*** If PT weight > **66kg (66Lb)** administer **Epi-Pen / Epinephrine Auto-Injector .3mg**

*** If PT weight < **66kg (66Lb)** administer **Epi-Pen JR / Epinephrine Auto-Injector .15mg**

Suspected Poisoning / Overdose

Activated Charcoal- Adult & Child: **1-2 gm/kg PO**

Diabetic Emergencies

If Blood Glucometry < **70mg/dL** AND Pt is conscious and able to speak and swallow administer **Oral Glucose 1 Tube**